



National
Qualifications
2019

2019 Mathematics
Higher Paper 1 (Non-calculator)
Finalised Marking Instructions

© Scottish Qualifications Authority 2019

These marking instructions have been prepared by examination teams for use by SQA appointed markers when marking external course assessments.

The information in this document may be reproduced in support of SQA qualifications only on a non-commercial basis. If it is reproduced, SQA must be clearly acknowledged as the source. If it is to be reproduced for any other purpose, written permission must be obtained from permissions@sqa.org.uk.



Marking instructions for each question

| Question | | Generic scheme | Illustrative scheme | Max mark |
|--|--|---|---|----------|
| 1. | | <ul style="list-style-type: none"> •¹ start to differentiate •² complete derivative and equate to 0 •³ factorise derivative •⁴ process cubic for x | <ul style="list-style-type: none"> •¹ $2x^3 \dots$ or $\dots - 6x^2$ •² $2x^3 - 6x^2 = 0$ •³ $2x^2(x-3)$ •⁴ 0 and 3 | 4 |
| Notes: | | | | |
| <ol style="list-style-type: none"> 1. •² is only available if '=0' appears at either •² or •³ stage, however see Candidate A. 2. Accept $2x^3 = 6x^2$ for •². 3. Accept $x^2(2x-6)$ for •³. 4. For candidates who divide by x or x^2 throughout see Candidate B. 5. •³ is available to candidates who factorise their derivative from •² as long as it is of equivalent difficulty. 6. $x=0$ and $x=3$ must be supported by valid working for •⁴ to be awarded. | | | | |
| Commonly Observed Responses: | | | | |
| Candidate A Stationary points when $\frac{dy}{dx} = 0$ $\frac{dy}{dx} = 2x^3 - 6x^2$ • ¹ ✓ • ² ✓ $\frac{dy}{dx} = 2x^2(x-3)$ • ³ ✓ $x=0$ and $x=3$ • ⁴ ✓ | | Candidate B $2x^3 - 6x^2 = 0$ • ¹ ✓ • ² ✓ $2x^3 = 6x^2$ • ³ ^ $x=3$ • ⁴ ✗ Dividing by x^2 is not valid as $x=0$ is a solution. | | |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|--|--|--|---|----------|
| 2. | | <ul style="list-style-type: none"> •¹ use discriminant •² apply condition and simplify •³ determine values of k | <ul style="list-style-type: none"> •¹ $(k-5)^2 - 4 \times 1 \times 1$ •² $k^2 - 10k + 21 = 0$ or $(k-5)^2 = 4$ •³ 3, 7 | 3 |
| Notes: | | | | |
| <p>1. Accept $(k-5)^2 - 4$ for •¹.</p> <p>2. Where candidates state an incorrect condition •² is not available. •³ is available for finding the roots of the quadratic. See Candidate B.</p> <p>3. Where x appears in any expression, no further marks are available.</p> | | | | |
| Commonly Observed Responses: | | | | |
| Candidate A | | Candidate B | | |
| For equal roots $b^2 - 4ac = 0$ | | For equal roots $b^2 - 4ac > 0$ | | |
| $(k-5)^2 - 4 \times 1 \times 1$ | | $(k-5)^2 - 4 \times 1 \times 1$ | | |
| $k^2 - 10k + 21$ | | $k^2 - 10k + 21 = 0$ or $(k-5)^2 = 4$ | | |
| $k = 3, 7$ | | $k = 3, 7$ | | |
| | | • ² ✘ | | |
| | | • ¹ ✓ | | |
| | | • ² ✓ | | |
| | | • ³ ✓ | | |
| | | • ³ ✓ 1 | | |
| Candidate C | | | | |
| $(k-5)^2 - 4 \times 1 \times 1 = 0$ | | | | |
| $k^2 - 10k = -21$ | | | | |
| $k = 3, 7$ | | | | |
| | | • ¹ ✓ | | |
| | | • ² ✓ | | |
| | | • ³ ✓ | | |
| No requirement for standard quadratic form | | | | |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|--|---|---|----------|
| 3. | | <ul style="list-style-type: none"> •¹ find radius of circle C_1 •² state equation of circle C_2 | <ul style="list-style-type: none"> •¹ 6 stated or implied by •² •² $(x-4)^2 + (y+2)^2 = 36$ | 2 |

Notes:

1. Accept $\sqrt{3^2 + 1^2 + 26} = 6$ or $\sqrt{-3^2 + -1^2 + 26} = 6$ for •¹.
2. Do not accept $\sqrt{-3^2 - 1^2 + 26} = 6$ for •¹.
3. Do not accept $(x-4)^2 + (y+2)^2 = 6^2$ for •².
4. For candidates whose working for $g^2 + f^2 - c$ does not arrive at a positive value, no marks are available. See Candidate A

Commonly Observed Responses:

Candidate A - 'fudging' negative values

$$\sqrt{3^2 + 1^2 - 26} = 4 \qquad \bullet^1 \times \bullet^2 \times$$

$$(x-4)^2 + (y+2)^2 = 16$$

| Question | | Generic scheme | Illustrative scheme | Max mark |
|--|-----|--|--|----------|
| 4. | (a) | <ul style="list-style-type: none"> •¹ interpret recurrence relation •² interpret recurrence relation •³ find m and c | <ul style="list-style-type: none"> •¹ $9 = 6m + c$ •² $11 = 9m + c$ •³ $m = \frac{2}{3}$ and $c = 5$ | 3 |
| Notes: | | | | |
| <p>1. Correct answer with no working award 0/3.</p> <p>2. Do not penalise $9 = m6 + c$ or $11 = m9 + c$ at •¹ and •².</p> <p>3. For candidates who state $m = \frac{2}{3}$, $c = 5$ and then verify that these values work for the given terms, award 2/3.</p> | | | | |
| Commonly Observed Responses: | | | | |
| | (b) | <ul style="list-style-type: none"> •⁴ calculate term | <ul style="list-style-type: none"> •⁴ $\frac{37}{3}$ or $12\frac{1}{3}$ | 1 |
| Notes: | | | | |
| <p>4. The answer in (b) must be consistent with the values found in (a).</p> <p>5. Accept $12.\dot{3}$ or $12.3\dots$ for •⁴. Do not accept a rounded answer.</p> | | | | |
| Commonly Observed Responses: | | | | |
| | | | | |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|-----|--|---|----------|
| 5. | (a) | <ul style="list-style-type: none"> •¹ find an appropriate vector eg \overrightarrow{AB} •² find a second vector eg \overrightarrow{BC} and compare •³ appropriate conclusion | <ul style="list-style-type: none"> •¹ eg $\overrightarrow{AB} = \begin{pmatrix} 3 \\ -6 \\ 3 \end{pmatrix}$ •² eg $\overrightarrow{BC} = \begin{pmatrix} 4 \\ -8 \\ 4 \end{pmatrix} \therefore \overrightarrow{AB} = \frac{3}{4}\overrightarrow{BC}$ •³ ... \Rightarrow AB is parallel to BC (common direction) and B is a common point \Rightarrow A, B and C are collinear. | 3 |

Notes:

- Do not penalise inconsistent vector notation (eg lack of arrows or brackets).
- Where •² is not awarded, if a candidate states that $\overrightarrow{AB} = \overrightarrow{BC}$, only •¹ is available.
- ³ can only be awarded if a candidate has stated 'parallel', 'common point' and 'collinear'.
- Candidates who state that 'points are parallel' or 'vectors are collinear' or 'parallel and share common point \Rightarrow collinear' do not gain •³. There must be reference to points A, B and C.
- Do not accept 'a, b and c are collinear' at •³.

Commonly Observed Responses:

Candidate A - missing labels

$$\begin{pmatrix} 3 \\ -6 \\ 3 \end{pmatrix} \quad \bullet^1 \wedge$$

$$\begin{pmatrix} 4 \\ -8 \\ 4 \end{pmatrix} \quad \therefore \overrightarrow{AB} = \frac{3}{4}\overrightarrow{BC} \quad \bullet^2 \checkmark 1$$

Missing labels at •2 is a repeated error

\Rightarrow AB is parallel to BC and B is a common point

\Rightarrow A, B and C are collinear $\bullet^3 \checkmark 1$

Candidate B

$$\overrightarrow{AB} = \begin{pmatrix} 3 \\ -6 \\ 3 \end{pmatrix} \quad \bullet^1 \checkmark$$

$$\overrightarrow{BC} = \begin{pmatrix} 4 \\ -8 \\ 4 \end{pmatrix}$$

$$\begin{pmatrix} 3 \\ -6 \\ 3 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix} \text{ and } \begin{pmatrix} 4 \\ -8 \\ 4 \end{pmatrix} = 4 \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix} \quad \bullet^2 \checkmark$$

$\therefore \overrightarrow{AB} = \frac{4}{3}\overrightarrow{BC}$

Ignore working subsequent to correct statement made on previous line.

\Rightarrow AB is parallel to BC and B is a common point

\Rightarrow A, B and C are collinear $\bullet^3 \checkmark$

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|--|----------------------------|---------------------|----------|
| (b) | | • ⁴ state ratio | • ⁴ 3:4 | 1 |

Notes:

- Answers in (b) must be consistent with the components of the vectors in (a) or the comparison of the vectors in (a). See Candidates C and D.
- In this case, the answer for •⁴ must be stated explicitly in part (b).
- The only acceptable variations for •⁴ must be related explicitly to AB and BC.
For $\frac{BC}{AB} = \frac{4}{3}$, $\frac{AB}{BC} = \frac{3}{4}$ or $BC:AB = 4:3$ stated in part (b) award •⁴. See Candidate E.
- Accept unitary ratios for •⁴, eg $\frac{3}{4}:1$ or $1:\frac{4}{3}$.
- Where a candidate states multiple ratios which are not equivalent, award 0/1.

Commonly Observed Responses:

Candidate C - using components of vectors

(a) $\vec{AB} = \begin{pmatrix} 3 \\ -6 \\ 3 \end{pmatrix}$ •¹ ✓

$\vec{BC} = \begin{pmatrix} 4 \\ -8 \\ 4 \end{pmatrix}$

$\vec{BC} = \frac{3}{4}\vec{AB}$ •² ✗

(b) 3:4 •⁴ ✓

Candidate D - using comparison of vectors

(a) $\vec{AB} = \begin{pmatrix} 3 \\ -6 \\ 3 \end{pmatrix}$ •¹ ✓

$\vec{BC} = \begin{pmatrix} 4 \\ -8 \\ 4 \end{pmatrix}$

$\vec{BC} = \frac{3}{4}\vec{AB}$ •² ✗

(b) 4:3 •⁴ ✓ 1

Candidate E - acceptable variation

$\frac{AB}{BC} = \frac{3}{4}$ •⁴ ✓

Ratio = 4:3

Ignore working subsequent to correct statement made on previous line.

Candidate F - trivial ratio

Ratio is 1:1 •⁴ ✓ 2

| Question | | Generic scheme | Illustrative scheme | Max mark |
|---|--|--|---|-----------------------------------|
| 6. | | <ul style="list-style-type: none"> •¹ write in differentiable form •² start to differentiate •³ complete differentiation | <ul style="list-style-type: none"> •¹ $(1-3x)^{-5}$ stated or implied by •² •² $-5(1-3x)^{-6} \dots$ •³ $\dots \times (-3)$ | 3 |
| Notes: | | | | |
| <p>1. Where candidates attempt to expand $(1-3x)^{-5}$, no further marks are available.</p> <p>2. •² is only available for differentiating an expression with a negative power.</p> | | | | |
| Commonly Observed Responses: | | | | |
| Candidate A | | | Candidate B | |
| $y = (1-3x)^{-5}$ | | • ¹ ✓ | $y = (1-3x)^{-5}$ | • ¹ ✓ |
| $\frac{dy}{dx} = -5(1-3x)^{-6} \times -3$ | | • ² ✓ • ³ ✓ | $\frac{dy}{dx} = -15(1-3x)^{-6}$ | • ² ✓ • ³ ✗ |
| $\frac{dy}{dx} = -15(1-3x)^{-6}$ | | | | |
| Candidate C | | | Candidate D - differentiating over two lines | |
| $y = (1-3x)^{-5}$ | | • ¹ ✓ | $y = (1-3x)^{-5}$ | • ¹ ✓ |
| . . . | | • ² ✓ • ³ ✗ | $\frac{dy}{dx} = -5(1-3x)^{-6}$ | • ² ✓ • ³ ^ |
| | | | $\frac{dy}{dx} = 15(1-3x)^{-6}$ | |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|--|---|--|----------|
| 7. | | <p>Method 1</p> <ul style="list-style-type: none"> •¹ use $m = \tan \theta$ •² find gradient of L •³ use property of perpendicular lines •⁴ determine equation of line | <p>Method 1</p> <ul style="list-style-type: none"> •¹ $m = \tan 30^\circ$ •² $\frac{1}{\sqrt{3}}$ •³ $-\sqrt{3}$ •⁴ $y = -\sqrt{3}x - 4$ | 4 |
| | | <p>Method 2</p> <ul style="list-style-type: none"> •¹ find angle perpendicular line makes with the positive direction of the x-axis. •² use $m = \tan \theta$ •³ find gradient of perpendicular line •⁴ determine equation of line | <p>Method 2</p> <ul style="list-style-type: none"> •¹ $30^\circ + 90^\circ = 120^\circ$ stated or implied by •² •² $m = \tan 120^\circ$ •³ $-\sqrt{3}$ •⁴ $y = -\sqrt{3}x - 4$ | |

Notes:

- In Method 1, where candidates make no reference to a trigonometric ratio or use an incorrect trigonometric ratio, •¹ and •² are unavailable.
In Method 2, where candidates use an incorrect trigonometric ratio •² and •³ are unavailable.
- Accept $y + 4 = -\sqrt{3}(x)$ at •⁴, but do not accept $y + 4 = -\sqrt{3}(x - 0)$.
- In Method 1, •⁴ is only available if the candidate has attempted to use a perpendicular gradient.

Commonly Observed Responses:

| | |
|--|---|
| <p>Candidate A</p> $m = \frac{1}{\sqrt{3}}$ (with or without diagram) • ¹ ^ • ² <input checked="" type="checkbox"/> 2 $m_{\perp} = -\sqrt{3}$ • ³ <input checked="" type="checkbox"/> 1 | <p>Candidate B</p> $m = \tan \theta$ (with or without diagram) • ¹ ^ $m = \frac{1}{\sqrt{3}}$ • ² <input checked="" type="checkbox"/> 1 |
| <p>Candidate C</p> $m = \tan \theta = 30$ • ¹ x $m = \frac{1}{\sqrt{3}}$ • ² <input checked="" type="checkbox"/> 1 | <p>Candidate D</p> $m = \tan^{-1} 30$ • ¹ x $m = \frac{1}{\sqrt{3}}$ • ² <input checked="" type="checkbox"/> 1 |
| <p>Candidate E</p> $\tan 30 = \frac{1}{\sqrt{3}}$ • ¹ ^ $m_{\perp} = -\sqrt{3}$ • ² <input checked="" type="checkbox"/> 1 • ³ <input checked="" type="checkbox"/> 1 | |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|--|-----|--|--|----------|
| 8. | (a) | • ¹ state integral | • ¹ $\int_{-1}^2 (-x^2 + x + 2) dx$ | 1 |
| Notes: | | | | |
| <p>1. Evidence for •¹ may be appear in part (b). However, where candidates make no attempt to answer part (a), •¹ is not available.</p> <p>2. •¹ is not available to candidates who omit the limits or 'dx'.</p> <p>3. •¹ is awarded for a candidates final expression for the area. However, accept $\int_{-1}^2 ((x^2 + 2x + 3) - (2x^2 + x + 1)) dx$ or $\int_{-1}^2 (x^2 + 2x + 3) dx - \int_{-1}^2 (2x^2 + x + 1) dx$ without further working.</p> <p>4. For $\int_{-1}^2 x^2 + 2x + 3 - 2x^2 + x + 1 dx$, see Candidates A and B.</p> | | | | |
| Commonly Observed Responses: | | | | |
| Candidate A | | Candidate B | | |
| (a) $\int_{-1}^2 x^2 + 2x + 3 - 2x^2 + x + 1 dx$ $\int_{-1}^2 (-x^2 + x + 2) dx$ • ¹ ✓ | | (a) $\int_{-1}^2 x^2 + 2x + 3 - 2x^2 + x + 1 dx$ (b) $\int_{-1}^2 (-x^2 + x + 2) dx$ • ¹ ✓ | | |
| Treat missing brackets as bad form as subsequent working is correct. | | • ¹ awarded in part (b) | | |
| Candidate C - error in simplification | | | | |
| (a) $\int_{-1}^2 (x^2 + 2x + 3) - (2x^2 + x + 1) dx$ $\int_{-1}^2 x^2 + x + 2 dx$ • ¹ ✗ | | | | |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|--|---|---|----------|
| (b) | | <ul style="list-style-type: none"> •² integrate expression from (a) •³ substitute limits •⁴ evaluate area | <ul style="list-style-type: none"> •² $-\frac{1}{3}x^3 + \frac{1}{2}x^2 + 2x$ •³ $\left(-\frac{1}{3}(2)^3 + \frac{1}{2}(2)^2 + 2(2)\right)$ $-\left(-\frac{1}{3}(-1)^3 + \frac{1}{2}(-1)^2 + 2(-1)\right)$ •⁴ $\frac{9}{2}$ | 3 |

Notes:

5. Where a candidate differentiates one or more terms at •² then •², •³ and •⁴ are unavailable.
6. Do not penalise the inclusion of '+c' or the continued appearance of the integral sign.
7. Candidates who substitute limits without integrating any term do not gain •³ or •⁴.
8. Where a candidate arrives at a negative value at •⁴ see Candidates D and E.

Commonly Observed Responses:

| | |
|--|---|
| <p>Candidate D</p> <p>Eg $\int_{-1}^2 (x^2 - x - 2) dx$</p> <p style="text-align: center;">⋮</p> <p style="text-align: center;">$= -\frac{9}{2} = \frac{9}{2}$ •⁴ ✘</p> <p>However...</p> <p style="text-align: center;">$= -\frac{9}{2}$, hence area is $\frac{9}{2}$. •⁴ ✔</p> | <p>Candidate E</p> <p>Eg $\int_{\frac{1}{2}}^{-1} (-x^2 + x + 2) dx$</p> <p style="text-align: center;">⋮</p> <p style="text-align: center;">$= -\frac{9}{2}$ cannot be negative so $\frac{9}{2}$ units² •⁴ ✘</p> <p>However...</p> <p style="text-align: center;">$= -\frac{9}{2}$, hence area is $\frac{9}{2}$. •⁴ ✔</p> |
| <p>Candidate F - not using expression from (a)</p> <p>(a) $\int_{-1}^2 x^2 + 2x + 3 dx$ •¹ ✘</p> <p>(b) $\int_{-1}^2 (x^2 + 2x + 3) - (2x^2 + x + 1) dx$</p> <p style="text-align: center;">$= \left[-\frac{1}{3}x^3 + \frac{1}{2}x^2 + 2x \right]_{-1}^2$ •² ✔ 2</p> <p style="text-align: center;">$= \left(-\frac{1}{3}(2)^3 + \frac{1}{2}(2)^2 + 2(2) \right)$</p> <p style="text-align: center;">$-\left(-\frac{1}{3}(-1)^3 + \frac{1}{2}(-1)^2 + 2(-1) \right)$ •³ ✔ 1</p> <p style="text-align: center;">$= \frac{9}{2}$ •⁴ ✔ 1</p> | |

| Question | | | Generic scheme | Illustrative scheme | Max mark |
|----------|-----|------|---|---|----------|
| 9. | (a) | (i) | • ¹ form an expression | • ¹ $p(2p+16)+(-2)(-3)+(4)(6)$ | 1 |
| | | (ii) | • ² equate scalar product to 0 • ³ factorise • ⁴ state values of p | • ² $p(2p+16)+(-2)(-3)+(4)(6)=0$ • ³ $2(p+5)(p+3)$ • ⁴ -5 and -3 | 3 |

Notes:

- Evidence for •¹ may appear in part (a)(ii).
- The appearance of ' $\mathbf{u} \cdot \mathbf{v} = 0$ ' alone is insufficient for •².
- For •² to be awarded '= 0' must appear at •² or •³.
- Do not penalise the absence of the common factor at •³.

Commonly Observed Responses:

| | |
|---|--|
| <p>Candidate A - incorrect expression at •²</p> <p>(i) $p(2p+16)+(-2)(-3)+(4)(6)$ •¹ ✓ $= 2p^2 + 16p + 30$ $= p^2 + 8p + 15$</p> <p>(ii) $p^2 + 8p + 15 = 0$ •² ✗ $(p+5)(p+3) = 0$ •³ ✓ 1 $p = -5, p = -3$ •⁴ ✓ 1</p> | <p>Candidate B - incorrect expression at •²</p> <p>(i) $p(2p+16)+(-2)(-3)+(4)(6)$ •¹ ✓ $= 2p^2 + 16p + 30$</p> <p>(ii) $p^2 + 8p + 15 = 0$ •² ✗ $(p+5)(p+3) = 0$ •³ ✓ 1 $p = -5, p = -3$ •⁴ ✓ 1</p> |
| <p>Candidate C - incorrect expression at •²</p> <p>$p(2p+16)+(-2)(-3)+(4)(6)$ •¹ ✓ $2p^2 + 16p + 24 = 0$ •² ✗ $2(p+6)(p+2)$ •³ ✓ 1 $p = -6, p = -2$ •⁴ ✓ 1</p> | <p>Candidate D</p> <p>(i) $\mathbf{u} \cdot \mathbf{v} = \begin{pmatrix} 2p^2 + 16p \\ 6 \\ 24 \end{pmatrix}$ •¹ ✗</p> <p>(ii) $p(2p+16)+6+24=0$ •² ✓ $2p^2 + 16p + 30 = 0$ $(p+5)(p+3) = 0$ •³ ✓ $p = -5, p = -3$ •⁴ ✓</p> |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|-----|---|--|----------|
| | (b) | <ul style="list-style-type: none"> •⁵ interpret relationship •⁶ determine value of p | <ul style="list-style-type: none"> •⁵ $3(p) = 2(2p + 16)$ or $3\mathbf{u} = 2\mathbf{v}$ or equivalent •⁶ -32 | 2 |

Notes:

Commonly Observed Responses:

Candidate E

For parallel vectors $\theta = 0^\circ$

Using $\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}| |\mathbf{v}| \cos \theta$

$$p(2p + 16) + (-2)(-3) + (4)(6) = \sqrt{p^2 + (-2)^2 + 4^2} \sqrt{(2p + 16)^2 + (-3)^2 + 6^2} \quad \bullet^5 \checkmark$$

$$p^2 + 64p + 1024 = 0$$

$$p = -32 \quad \bullet^6 \checkmark$$

| Question | | Generic scheme | Illustrative scheme | Max mark |
|-------------------------------------|-----|--------------------------------------|---------------------|----------|
| 10. | (a) | • ¹ identify value of a | • ¹ 3 | 1 |
| Notes: | | | | |
| | | | | |
| Commonly Observed Responses: | | | | |
| | | | | |
| | (b) | • ² identify value of k | • ² -2 | 1 |
| Notes: | | | | |
| | | | | |
| Commonly Observed Responses: | | | | |
| | | | | |

| Question | Generic scheme | Illustrative scheme | Max mark |
|----------|--|---|----------|
| 11. | <ul style="list-style-type: none"> •¹ start to integrate •² complete integration •³ substitute limits •⁴ evaluate integral | <ul style="list-style-type: none"> •¹ $\sin\left(3x - \frac{\pi}{6}\right) \dots$ •² $\dots \times \frac{1}{3}$ •³ $\left(\frac{1}{3} \sin\left(3 \times \frac{\pi}{9} - \frac{\pi}{6}\right)\right)$ $-\left(\frac{1}{3} \sin\left(3 \times 0 - \frac{\pi}{6}\right)\right)$ •⁴ $\frac{1}{3}$ | 4 |

Notes:

1. Where candidates make no attempt to integrate or start to integrate individual terms within the bracket or use another invalid approach eg $\sin\left(3x - \frac{\pi}{6}\right)^2$ or $\int \cos(3x) - \cos\left(\frac{\pi}{6}\right) dx$, award 0/4.
2. Do not penalise the inclusion of '+c' or the continued appearance of the integral sign after •¹.
3. Candidates who work in degrees from the start cannot gain •¹. However, •², •³ and •⁴ are still available.
4. •¹ may be awarded for the appearance of $\sin\left(3x - \frac{\pi}{6}\right)$ in the first line of working, however see Candidates B and D.
5. •⁴ is only available where candidates have considered both limits within a trigonometric function.
6. Where candidates use a mixture of degrees and radians, •³ is not awarded. However, •⁴ is still available.

Commonly Observed Responses:

| | |
|--|---|
| <p>Candidate A - using addition formula</p> $\int_0^{\frac{\pi}{9}} \left(\cos 3x \cos \frac{\pi}{6} + \sin 3x \sin \frac{\pi}{6} \right) dx$ $= \frac{1}{3} \sin 3x \times \frac{\sqrt{3}}{2} \dots$ $\dots - \frac{1}{3} \cos 3x \times \frac{1}{2}$ <p>•¹ ✓ •² ✓</p> | <p>Candidate B - integrated over two lines</p> $\int_0^{\frac{\pi}{9}} \left(\cos \left(3x - \frac{\pi}{6} \right) \right) dx$ $= \sin \left(3x - \frac{\pi}{6} \right)$ $= \frac{1}{3} \sin \left(3x - \frac{\pi}{6} \right)$ <p>•¹ ✓ •² ✗</p> |
| <p>Candidate C - integrated in part</p> $3 \sin \left(3x - \frac{\pi}{6} \right)$ $3 \sin \left(3 \times \frac{\pi}{9} - \frac{\pi}{6} \right) - 3 \sin \left(0 - \frac{\pi}{6} \right)$ 3 <p>•¹ ✓ •² ✗</p> <p>•³ ✓ 1</p> <p>•⁴ ✓ 1</p> | <p>Candidate D - integrated in part</p> $-\frac{1}{3} \sin \left(3x - \frac{\pi}{6} \right)$ $-\frac{1}{3} \sin \left(3 \times \frac{\pi}{9} - \frac{\pi}{6} \right) + \frac{1}{3} \sin \left(0 - \frac{\pi}{6} \right)$ $-\frac{1}{3}$ <p>•¹ ✗ •² ✓</p> <p>•³ ✓ 1</p> <p>•⁴ ✓ 1</p> |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|---|-----|---|--|----------|
| 12. | (a) | <ul style="list-style-type: none"> •¹ interpret notation •² state expression for $f(g(x))$ | <ul style="list-style-type: none"> •¹ $f(5-x)$ or $\frac{1}{\sqrt{g(x)}}$ •² $\frac{1}{\sqrt{5-x}}$ | 2 |
| Notes: | | | | |
| 1. For $\frac{1}{\sqrt{5-x}}$ without working, award both • ¹ and • ² . | | | | |
| Commonly Observed Responses: | | | | |
| Candidate A | | | | |
| $5 - \frac{1}{\sqrt{x}}$ | | • ¹ ✗ • ² <input checked="" type="checkbox"/> 1 | | |
| | (b) | • ³ state range | • ³ $x \geq 5$ | 1 |
| Notes: | | | | |
| 2. Answer at • ³ must be consistent with expression at • ² . | | | | |
| 3. For candidates who interpret $g(f(x))$ as $f(g(x))$, do not award • ³ . | | | | |
| Commonly Observed Responses: | | | | |
| Candidate B | | | | |
| $5 - \frac{1}{\sqrt{x}}$ | | • ¹ ✗ • ² <input checked="" type="checkbox"/> 1 | | |
| $x \leq 0$ | | • ³ ✗ | | |

| Question | | | Generic scheme | Illustrative scheme | Max mark |
|----------|-----|------|-----------------------------------|--------------------------------------|----------|
| 13. | (a) | (i) | • ¹ determine $\cos p$ | • ¹ $\frac{2}{\sqrt{5}}$ | 1 |
| | | (ii) | • ² determine $\cos q$ | • ² $\frac{3}{\sqrt{10}}$ | 1 |

Notes:

1. Where candidates do not simplify the perfect squares see Candidates A and B.

Commonly Observed Responses:

Candidate A - no evidence of simplification

$$\cos p = \frac{\sqrt{4}}{\sqrt{5}} \quad \bullet^1 \times$$

$$\cos q = \frac{\sqrt{9}}{\sqrt{10}} \quad \bullet^2 \boxed{1}$$

Repeated error not penalised twice

Candidate B - simplification in part (b)

$$(a) \cos p = \frac{\sqrt{4}}{\sqrt{5}} \cos q = \frac{\sqrt{9}}{\sqrt{10}} \quad \bullet^1 \checkmark$$

$$\vdots$$

$$(b) \sin(p+q) = \frac{5}{\dots} \quad \bullet^2 \checkmark$$

Roots have been simplified in (b)

| Question | | | Generic scheme | Illustrative scheme | mark |
|----------|-----|--|---|---|------|
| | (b) | | • ³ select appropriate formula and express in terms of p and q • ⁴ substitute into addition formula • ⁵ evaluate $\sin(p+q)$ | • ³ $\sin p \cos q + \cos p \sin q$ • ⁴ $\frac{1}{\sqrt{5}} \times \frac{3}{\sqrt{10}} + \frac{2}{\sqrt{5}} \times \frac{1}{\sqrt{10}}$ • ⁵ $\frac{1}{\sqrt{2}}$ | 3 |

Notes:

2. Award •³ for candidates who write $\sin\left(\frac{1}{\sqrt{5}}\right) \times \cos\left(\frac{3}{\sqrt{10}}\right) + \cos\left(\frac{2}{\sqrt{5}}\right) \times \sin\left(\frac{1}{\sqrt{10}}\right)$. •⁴ and •⁵ are unavailable.

3. For any attempt to use $\sin(p+q) = \sin p + \sin q$, •⁴ and •⁵ are unavailable.

4. At •⁵, accept answers such as $\frac{5}{\sqrt{50}}$ or $\frac{5}{5\sqrt{2}}$ but not $\frac{5}{\sqrt{5} \times \sqrt{10}}$.

5. At •⁵, the answer must be given as a single fraction.

6. Do not penalise trigonometric ratios which are less than -1 or greater than 1 .

Commonly Observed Responses:

| Question | | Generic scheme | Illustrative scheme | Max mark | | |
|---|--|---|---|----------|---|--|
| 14. | (a) | <ul style="list-style-type: none"> •¹ apply $m \log_n x = \log_n x^m$ •² apply •³ evaluate logarithm | <ul style="list-style-type: none"> •¹ ...$\log_{10} 5^2$ stated or implied by •² •² $\log_{10}(4 \times 5^2)$ •³ 2 | 3 | | |
| Notes: | | | | | | |
| <p>1. Each line of working must be equivalent to the line above within a valid strategy, however see Candidate A.</p> <p>2. Do not penalise the omission of the base of the logarithm at •¹ or •².</p> <p>3. Correct answer with no working, award 0/3.</p> | | | | | | |
| Commonly Observed Responses: | | | | | | |
| <table border="0" style="width: 100%;"> <tr> <td style="width: 50%; vertical-align: top;"> Candidate A $2 \log_{10}(4 \times 5)$ $2 \log_{10}(20)$ $\log_{10}(20)^2$ </td> <td style="width: 50%; vertical-align: top; text-align: center;"> <ul style="list-style-type: none"> •² ✘ •¹ 1 •³ ^ </td> </tr> </table> | | | | | Candidate A $2 \log_{10}(4 \times 5)$ $2 \log_{10}(20)$ $\log_{10}(20)^2$ | <ul style="list-style-type: none"> •² ✘ •¹ 1 •³ ^ |
| Candidate A $2 \log_{10}(4 \times 5)$ $2 \log_{10}(20)$ $\log_{10}(20)^2$ | <ul style="list-style-type: none"> •² ✘ •¹ 1 •³ ^ | | | | | |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|--|-----|---|---|---|
| | (b) | <p>Method 1</p> <ul style="list-style-type: none"> •⁴ apply $\log_a x - \log_a y = \log_a \frac{x}{y}$ •⁵ express in exponential form •⁶ solve for x | <p>Method 1</p> <ul style="list-style-type: none"> •⁴ $\log_2 \frac{7x-2}{3} = \dots$ •⁵ $\frac{7x-2}{3} = 2^5$ •⁶ 14 | 3 |
| | | <p>Method 2</p> <ul style="list-style-type: none"> •⁴ apply $m \log_n x = \log_n x^m$ •⁵ simplify •⁶ solve for x | <p>Method 2</p> <ul style="list-style-type: none"> •⁴ $\dots = \log_2 2^5$ •⁵ eg $\log_2 \frac{7x-2}{3} = \dots$ or $\log_2 (7x-2) = \log_2 (3 \times 2^5)$ •⁶ 14 | |
| Notes: | | | | |
| 4. • ⁶ is only awarded if each line of working is equivalent to the line above within a valid strategy. | | | | |
| Commonly Observed Responses: | | | | |
| Candidate A - invalid working leading to solution | | Candidate B - invalid working leading to solution | | |
| $\log_2 \frac{7x-2}{3} = \log_2 5^2$ $x = 11$ | | <ul style="list-style-type: none"> •⁴ ✓ •⁵ ✗ •⁶ <input checked="" type="checkbox"/> 2 | $\log_2 \frac{7x-2}{3} = \log_2 5 \times 2$ $x = \frac{32}{7}$ | <ul style="list-style-type: none"> •⁴ ✓ •⁵ ✗ •⁶ <input checked="" type="checkbox"/> 2 |
| Candidate C | | Candidate D | | |
| $\log_2 \left(\frac{7x-2}{3} \right) = 5 \log_2 2$ $\log_2 \frac{7x-2}{3} - \frac{2}{3} = \log_2 2^5$ | | <ul style="list-style-type: none"> •⁵ ✓ •⁴ ✓ | $\log_2 (7x-2) - \log_2 3 = \log_2 2^5$ $\log_2 \left(\frac{7x-2}{3} \right) = \log_2 25$ | <ul style="list-style-type: none"> •⁴ ✓ •⁵ ✓ |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|-----|---|---|----------|
| 15. | (a) | <ul style="list-style-type: none"> •¹ substitute appropriate double angle formula •² factorise •³ solve for $\cos x^\circ$ and $\sin x^\circ$ •⁴ solve for x | <ul style="list-style-type: none"> •¹ $2 \sin x^\circ \cos x^\circ + 6 \cos x^\circ = 0$ •² $2 \cos x^\circ (\sin x^\circ + 3) = 0$ •³ $\cos x^\circ = 0$ •⁴ $\sin x^\circ = -3$ •⁴ $x = 90, 270$ 'no solutions' | 4 |

Notes:

1. Do not penalise the absence of '=0' at •¹ and •².
2. Do not penalise the absence of '2' as a common factor at •².
3. Do not penalise the omission of degree signs.
4. Candidates who leave their answer in radians do not gain •⁴ (if marking horizontally) or •³ (if marking vertically).
5. •⁴ is only available if one of the equations at •³ has no solution.
6. Accept ~~$\sin x^\circ = -3$~~ at •⁴.

Commonly Observed Responses:

| | | |
|---|--|--|
| <p>Candidate A</p> $2 \sin x \cos x = -6 \cos x$ $2 \sin x = -6$ $\sin x = -3$ | <ul style="list-style-type: none"> •¹ ✓ •² ^ •³ ^ •⁴ ✓ 1 | <p>Candidate B - insufficient evidence for •³</p> $2 \sin x^\circ \cos x^\circ + 6 \cos x^\circ = 0$ • ¹ ✓ $2 \cos x^\circ (\sin x^\circ + 3) = 0$ • ² ✓ $2 \cos x^\circ = 0, \sin x^\circ = -3$ • ³ ^ • ⁴ ^ However, $x = 90, 270, \text{'no solutions'}$ • ³ ✓ • ⁴ ✓ |
|---|--|--|

| | | | |
|-----|--------------------------------|----------------------------------|---|
| (b) | • ⁵ state solutions | • ⁵ 45, 135, 225, 315 | 1 |
|-----|--------------------------------|----------------------------------|---|

Notes:

Commonly Observed Responses:

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|-----|--|---|----------|
| 16. | (a) | <ul style="list-style-type: none"> •¹ identify centre •² apply distance formula and demonstrate result | <ul style="list-style-type: none"> •¹ (1, -2) stated or implied by •² •² $\sqrt{(4-1)^2 + (k - (-2))^2}$ leading to $\sqrt{k^2 + 4k + 13}$ | 2 |

Notes:

1. Beware of candidates who 'fudge' their working between •¹ and •².

Commonly Observed Responses:

| | | | | |
|--|-----|---|---|---|
| | (b) | <ul style="list-style-type: none"> •³ interpret information •⁴ express inequality in standard quadratic form •⁵ determine zeros of quadratic expression •⁶ state range with justification | <ul style="list-style-type: none"> •³ $\sqrt{k^2 + 4k + 13} > 5$ •⁴ $k^2 + 4k - 12 > 0$ •⁵ -6, 2 •⁶ $k < -6, k > 2$ with eg sketch or table of signs | 4 |
|--|-----|---|---|---|

Notes:

2. Where a candidate has used an incorrect expression from part (a), •³ is not available. However, •⁴, •⁵ and •⁶ are still available for dealing with an expression of equivalent difficulty.
3. Candidates who do not work with an inequation from the outset lose •³, •⁴ and •⁶. However, •⁵ is still available. See Candidate A.

Commonly Observed Responses:

| | |
|---|--|
| <p>Candidate A</p> <p>$\sqrt{k^2 + 4k + 13} = 5$</p> <p>$k^2 + 4k - 12 = 0$</p> <p>$k = -6, k = 2$</p> <p>For P to lie outside the circle</p> <p>$k < -6, k > 2$</p> | <ul style="list-style-type: none"> •³ ✗ •⁴ ✗ •⁵ ✓ •⁶ ✗ |
|---|--|

| Question | | Generic scheme | Illustrative scheme | Max mark |
|---|-----|--|--|----------|
| 17. | (a) | <ul style="list-style-type: none"> •¹ expand brackets •² use double angle formula for sin •³ use trigonometric identity and express in required form | <ul style="list-style-type: none"> •¹ $\sin^2 x - \sin x \cos x$ $-\sin x \cos x + \cos^2 x$ •² $\dots - \sin 2x \dots$ •³ $1 - \sin 2x$ | 3 |
| Notes: | | | | |
| 1. For correct answer with no working award 0/3. | | | | |
| Commonly Observed Responses: | | | | |
| Candidate A - incorrect notation $\sin x^2 - 2 \sin x \cos x + \cos x^2$ • ¹ ✗ $1 - \sin 2x$ • ² ✓ • ³ ✗ | | | | |
| | (b) | <ul style="list-style-type: none"> •⁴ link to (a) and integrate one term •⁵ complete integration | <ul style="list-style-type: none"> •⁴ eg $\int (1 - \sin 2x) dx = x \dots$ •⁵ $x + \frac{1}{2} \cos 2x + c$ | 2 |
| Notes: | | | | |
| 2. • ⁴ and • ⁵ can only be awarded if the integrand is of the form $p + q \sin rx$. | | | | |
| 3. Where the statement for • ³ appears with no relevant working, • ⁴ and • ⁵ are not available. | | | | |
| Commonly Observed Responses: | | | | |
| | | | | |

[END OF MARKING INSTRUCTIONS]



National
Qualifications
2019

2019 Mathematics

Higher Paper 2

Finalised Marking Instructions

© Scottish Qualifications Authority 2019

These marking instructions have been prepared by examination teams for use by SQA appointed markers when marking external course assessments.

The information in this document may be reproduced in support of SQA qualifications only on a non-commercial basis. If it is reproduced, SQA must be clearly acknowledged as the source. If it is to be reproduced for any other purpose, written permission must be obtained from permissions@sqa.org.uk.



©

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|-----|--|---|----------|
| 1. | (a) | <ul style="list-style-type: none"> •¹ calculate the midpoint of AC •² calculate the gradient of BD •³ determine equation of BD | <ul style="list-style-type: none"> •¹ $(-4, -3)$ •² $-\frac{1}{3}$ •³ $3y = -x - 13$ | 3 |

Notes:

1. •² is only available to candidates who use a midpoint to find a gradient.
2. •³ is only available as a consequence of using the midpoint of AC and the point B.
3. At •³ accept any arrangement of a candidate's equation where constant terms have been simplified.
4. •³ is not available as a consequence of using a perpendicular gradient.

Commonly Observed Responses:

| | |
|---|--|
| <p>Candidate A - Perpendicular Bisector of AC Midpoint_{AC} $(-4, -3)$ •¹ ✓ $m_{AC} = 9 \Rightarrow m_{\perp} = -\frac{1}{9}$ •² ✗ $9y + x + 31 = 0$ •³ ✓ 2 For other perpendicular bisectors award 0/3</p> | <p>Candidate B - Altitude through B $m_{AC} = 9$ •¹ ^ $m_{\perp} = -\frac{1}{9}$ •² ✗ $9y + x = -61$ •³ ✓ 2</p> |
| <p>Candidate C - Median through A Midpoint_{BC} $(4, -1)$ •¹ ✗ $m_{AM} = \frac{11}{9}$ •² ✓ 1 $9y - 11x + 53 = 0$ •³ ✓ 2</p> | <p>Candidate D - Median through C Midpoint_{AB} $(3, -10)$ •¹ ✗ $m_{CM} = -\frac{8}{3}$ •² ✓ 1 $3y + 8x + 6 = 0$ •³ ✓ 2</p> |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|---|-----|---|--|----------|
| | (b) | <ul style="list-style-type: none"> •⁴ calculate gradient of BC •⁵ use property of perpendicular lines •⁶ determine equation of AE | <ul style="list-style-type: none"> •⁴ -1 •⁵ 1 •⁶ $y = x - 7$ | 3 |
| Notes: | | | | |
| <p>5. •⁶ is only available to candidates who find and use a perpendicular gradient.</p> <p>6. At •⁶ accept any arrangement of a candidate's equation where constant terms have been simplified.</p> | | | | |
| Commonly Observed Responses: | | | | |
| <p>Candidate E Correct gradient from incorrect substitution</p> $m_{BC} = \frac{-3 - 11}{6 + 8} = -1 \quad \bullet^4 \times$ $m_{AE} = 1 \quad \bullet^5 \boxed{\checkmark 1}$ $y = x - 7 \quad \bullet^6 \boxed{\checkmark 1}$ | | | | |
| | (c) | <ul style="list-style-type: none"> •⁷ find x or y coordinate •⁸ find remaining coordinate of the point of intersection | <ul style="list-style-type: none"> •⁷ $x = 2$ or $y = -5$ •⁸ $y = -5$ or $x = 2$ | 2 |
| Notes: | | | | |
| 7. For $(2, -5)$ with no working, award 0/2. | | | | |
| Commonly Observed Responses: | | | | |
| | | | | |

| Question | Generic scheme | Illustrative scheme | Max mark |
|----------|--|---|----------|
| 2. | <ul style="list-style-type: none"> •¹ express $6\sqrt{x}$ in integrable form •² integrate first term •³ integrate second term •⁴ complete integration | <ul style="list-style-type: none"> •¹ $6x^{\frac{1}{2}}$ •² $\frac{6x^{\frac{3}{2}}}{\frac{3}{2}} \dots$ •³ $\dots - \frac{4x^{-2}}{-2} \dots$ •⁴ $4x^{\frac{3}{2}} + 2x^{-2} + 5x + c$ | 4 |

Notes:

1. •² is only available for integrating a term with a fractional index.
2. All coefficients must be simplified at •⁴ stage for •⁴ to be awarded.
3. Do not penalise the appearance of an integral sign throughout.
4. Do not penalise the omission of '+c' at •² and •³.

Commonly Observed Responses:

Candidate A

$$\int \left(6x^{\frac{1}{2}} - 4x^{-3} + 5 \right) dx \quad \bullet^1 \checkmark$$

$$= \frac{6x^{\frac{3}{2}}}{\frac{3}{2}} - \frac{4x^{-2}}{-2} + 5x + c \quad \bullet^2 \checkmark \quad \bullet^3 \checkmark$$


$$= \frac{12}{3} x^{\frac{3}{2}} + 2x^{-2} + 5x + c$$

$$= 4\sqrt{x^3} + \frac{2}{\sqrt{x}} + 5x + c \quad \bullet^4 \times$$

•⁴ cannot be awarded over two lines of working

| Question | | Generic scheme | Illustrative scheme | Max mark |
|--|-----|--|---|----------|
| 3. | (a) | • ¹ identify pathway | • ¹ $-\mathbf{p} + \mathbf{r}$ | 1 |
| Notes: | | | | |
| 1. Accept $-\mathbf{P} + \mathbf{R}$ for • ¹ . | | | | |
| Commonly Observed Responses: | | | | |
| | (b) | • ² state an appropriate pathway • ³ express pathway in terms of \mathbf{p} , \mathbf{q} and \mathbf{r} | • ² eg $\overrightarrow{EB} + \overrightarrow{BF}$ stated or implied by • ³ • ³ $\mathbf{p} - \mathbf{r} + \frac{3}{4}\mathbf{q}$ or equivalent | 2 |
| Notes: | | | | |
| 2. • ³ can only be awarded for a vector expressed in terms of all three of \mathbf{p} , \mathbf{q} and \mathbf{r} . | | | | |
| Commonly Observed Responses: | | | | |
| Candidate A - incorrect expression in \mathbf{p} , \mathbf{q} and \mathbf{r} and no pathway stated $\mathbf{p} - \mathbf{r} \dots$ | | Candidate B - incorrect expression in \mathbf{p} , \mathbf{q} and \mathbf{r} and no pathway stated $\dots + \frac{3}{4}\mathbf{q}$ or $\dots + \mathbf{q} - \frac{1}{4}\mathbf{q}$ | | |
| Award 1/2 | | Award 1/2 | | |

| Question | | Generic scheme | | Illustrative scheme | Max mark |
|---|-----|----------------|--|--|----------|
| 4. | (a) | | • ¹ state values of a and b | • ¹ $a = 0.973, b = 30$ | 1 |
| Notes: | | | | | |
| 1. Accept $u_{n+1} = 0.973u_n + 30$ for • ¹ . | | | | | |
| Commonly Observed Responses: | | | | | |
| | (b) | (i) | • ² communicate condition for limit to exist | • ² a limit exists as the recurrence relation is linear and $-1 < 0.973 < 1$ | 1 |
| | | (ii) | • ³ know how to find limit • ⁴ process limit and state estimated population | • ³ $L = 0.973L + 30$ or $L = \frac{30}{1 - 0.973}$ • ⁴ 1100 | 2 |
| Notes: | | | | | |
| 2. For • ² accept: $-1 < 0.973 < 1$ or $ 0.973 < 1$ or $0 < 0.973 < 1$ with no further comment; or statements such as “0.973 lies between -1 and 1”; or $-1 < a < 1$ (as a is previously defined). 3. • ² is not available for: $-1 \leq 0.973 \leq 1$ or $0.973 < 1$; or statements such as “it is between -1 and 1” 4. Do not accept $L = \frac{b}{1-a}$ with no further working for • ³ . 5. For $L = 1100$ with no working award • ³ and • ⁴ . | | | | | |
| Commonly Observed Responses: | | | | | |
| Candidate A - no rounding required | | | Candidate B - correct rounding | | |
| $u_{n+1} = 0.97u_n + 30$ | | | $u_{n+1} = 0.027u_n + 30$ | | |
| ⋮ | | | ⋮ | | |
| $L = \frac{30}{1 - 0.97}$ | | | $L = \frac{30}{1 - 0.027}$ | | |
| $L = 1000$ | | | $L = 0$ | | |
| • ¹ ✗ | | | • ¹ ✗ | | |
| • ³ ✓ 1 | | | • ³ ✓ 1 | | |
| • ⁴ ✓ 2 | | | • ⁴ ✓ 1 | | |
| Candidate C - no valid limit | | | | | |
| $u_{n+1} = 2.7u_n + 30$ | | | • ¹ ✗ | | |
| A limit does not exist as $2.7 > 1$ | | | • ² ✗ | | |
| $L = \frac{30}{1 - 2.7}$ | | | • ³ ✓ 1 | | |
| $L = 0$ | | | • ⁴ ✗ | | |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|--|--|---|---|----------|
| 5. | | <ul style="list-style-type: none"> •¹ identify shape and roots •² interpret shape | <ul style="list-style-type: none"> •¹ parabola with roots at -2 and 4 •² parabola with a minimum turning point at $x = 1$  | 2 |
| Notes: | | | | |
| 1. • ¹ and • ² are only available for attempting to draw a 'parabola'. | | | | |
| Commonly Observed Responses: | | | | |
| | | | | |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|-----|---|--|----------|
| 6. | (a) | <ul style="list-style-type: none"> •¹ use compound angle formula •² compare coefficients •³ process for k •⁴ process for a and express in required form | <ul style="list-style-type: none"> •¹ $k \cos x^\circ \cos a^\circ - k \sin x^\circ \sin a^\circ$ stated explicitly •² $k \cos a^\circ = 2, k \sin a^\circ = 3$ stated explicitly •³ $\sqrt{13}$ •⁴ $\sqrt{13} \cos(x + 56 \cdot 3 \dots)^\circ$ | 4 |

Notes:

1. Accept $k(\cos x^\circ \cos a^\circ - \sin x^\circ \sin a^\circ)$ for •¹.
Treat $k \cos x^\circ \cos a^\circ - \sin x^\circ \sin a^\circ$ as bad form only if the equations at the •² stage both contain k .
2. Do not penalise the omission of degree signs.
3. $\sqrt{13} \cos x^\circ \cos a^\circ - \sqrt{13} \sin x^\circ \sin a^\circ$ or $\sqrt{13}(\cos x^\circ \cos a^\circ - \sin x^\circ \sin a^\circ)$ is acceptable for •¹ and •³.
4. •² is not available for $k \cos x^\circ = 2, k \sin x^\circ = 3$, however •⁴ may still be gained. See Candidate F.
5. Accept $k \cos a^\circ = 2, -k \sin a^\circ = -3$ for •².
6. •³ is only available for a single value of $k, k > 0$.
7. •⁴ is not available for a value of a given in radians.
8. Accept values of a which round to 56.
9. Candidates may use any form of the wave function for •¹, •² and •³.
However, •⁴ is only available if the wave is interpreted in the form $k \cos(x + a)^\circ$.
10. Evidence for •⁴ may not appear until part (b).

Commonly Observed Responses:

| Candidate A | Candidate B | Candidate C |
|--|---|--|
| $\sqrt{13} \cos a^\circ = 2$ $\sqrt{13} \sin a^\circ = 3$ $\tan a^\circ = \frac{3}{2}$ $a = 56 \cdot 3$ $\sqrt{13} \cos(x + 56 \cdot 3)^\circ$ | $k \cos x^\circ \cos a^\circ - k \sin x^\circ \sin a^\circ$ $\cos a^\circ = 2$ $\sin a^\circ = 3$ $\tan a^\circ = \frac{3}{2}$ $a = 56 \cdot 3$ $\sqrt{13} \cos(x + 56 \cdot 3)^\circ$ | $\cos x^\circ \cos a^\circ - \sin x^\circ \sin a^\circ$ $\cos a^\circ = 2$ $\sin a^\circ = 3$ $k = \sqrt{13}$ $\tan a^\circ = \frac{3}{2}$ $a = 56 \cdot 3$ $\sqrt{13} \cos(x + 56 \cdot 3)^\circ$ |
| <ul style="list-style-type: none"> •¹ ^ •² ✓ •³ ✓ •⁴ ✓ | <ul style="list-style-type: none"> •¹ ✓ •² ✗ •³ ✓ •⁴ ✗ | <ul style="list-style-type: none"> •¹ ✗ •² ✓ 2 •³ ✓ •⁴ ✗ |

| Question | Generic scheme | Illustrative scheme | Max mark |
|---|---|--|----------|
| <p>Candidate D - errors at \bullet^2 $k \cos x^\circ \cos a^\circ - k \sin x^\circ \sin a^\circ$ $\bullet^1 \checkmark$ $k \cos a^\circ = 3$ $k \sin a^\circ = 2$ $\bullet^2 \times$ $\tan a^\circ = \frac{2}{3}$ $a = 33.7$ $\sqrt{13} \cos(x+33.7)^\circ$ $\bullet^3 \checkmark$ $\bullet^4 \checkmark 1$</p> | <p>Candidate E - errors at \bullet^2 $k \cos x^\circ \cos a^\circ - k \sin x^\circ \sin a^\circ$ $\bullet^1 \checkmark$ $k \cos a^\circ = 2$ $k \sin a^\circ = -3$ $\bullet^2 \times$ $\tan a^\circ = -\frac{3}{2}$ $a = 303.7$ $\sqrt{13} \cos(x+303.7)^\circ$ $\bullet^3 \checkmark$ $\bullet^4 \checkmark 1$</p> | <p>Candidate F - use of x $k \cos x^\circ \cos a^\circ - k \sin x^\circ \sin a^\circ$ $\bullet^1 \checkmark$ $k \cos x^\circ = 2$ $k \sin x^\circ = 3$ $\bullet^2 \times$ $\tan a^\circ = \frac{3}{2}$ $x = 56.3$ $\sqrt{13} \cos(x+56.3)^\circ$ $\bullet^3 \checkmark$ $\bullet^4 \checkmark 1$</p> | |
| <p>Candidate G $k \cos A \cos B - k \sin A \sin B$ $\bullet^1 \times$ $k \cos A^\circ = 2$ $k \sin A^\circ = 3$ $\bullet^2 \times$ $\tan A^\circ = \frac{3}{2}$ $a = 56.3$ $\sqrt{13} \cos(x+56.3)^\circ$ $\bullet^3 \checkmark$ $\bullet^4 \checkmark 1$</p> <p>Unclear at this stage whether A relates to a or to x.</p> | | | |
| (b) | <ul style="list-style-type: none"> \bullet^5 link to (a) \bullet^6 solve for $x+a$ \bullet^7 solve for x | $\bullet^5 \sqrt{13} \cos(x+56.3\dots)^\circ = 3$ $\bullet^6 33.69\dots(393.69\dots)$ $\bullet^7 326.31\dots$ $\bullet^7 337.38\dots$ 270 | 3 |
| Notes: | | | |
| 11. Do not penalise working which rounds to 34, 326, 394 leading to 270 and 337. | | | |
| Commonly Observed Responses: | | | |
| | | | |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|-----|--|--|----------|
| 7. | (a) | <p>Method 1</p> <ul style="list-style-type: none"> •¹ identify common factor •² complete the square •³ process for r and write in required form | <p>Method 1</p> <ul style="list-style-type: none"> •¹ $-6(x^2 - 4x...)$ stated or implied by •² •² $-6(x-2)^2 ...$ •³ $-6(x-2)^2 - 1$ | 3 |
| | | <p>Method 2</p> <ul style="list-style-type: none"> •¹ expand completed square form •² equate coefficients •³ process for q and r and write in required form | <p>Method 2</p> <ul style="list-style-type: none"> •¹ $px^2 + 2pqx + pq^2 + r$ •² $p = -6, 2pq = 24, pq^2 + r = -25$ •³ $-6(x-2)^2 - 1$ | |

Notes:

- $-6(x-2)^2 - 1$ with no working gains •¹ and •² only. However, see Candidate E.
- ³ is not available in cases where $p > 0$.

Commonly Observed Responses:

| | |
|--|--|
| <p>Candidate A</p> $-6(x^2 - 4) - 25$ $-6((x-2)^2 - 4) - 25$ • ¹ ✓ • ² ✓ $-6(x-2)^2 - 1$ • ³ ✓ See the exception to general marking principle (h) | <p>Candidate B</p> $px^2 + 2pqx + pq^2 + r$ • ¹ ✓ $p = -6, 2pq = 24, pq^2 + r = -25$ • ² ✓ $q = -2, r = -1$ • ³ ^ <div style="border: 1px solid black; border-radius: 15px; padding: 5px; display: inline-block; margin-top: 10px;"> •³ is lost as answer is not in completed square form </div> |
| <p>Candidate C</p> $-6(x^2 + 24x) - 25$ • ¹ ✗ $-6((x+12)^2 - 144) - 25$ • ² ✓ 1 $-6(x+12)^2 + 839$ • ³ ✓ 1 | <p>Candidate D</p> $-6((x+12)^2 - 144) - 25$ • ¹ ^ • ² ✗ $-6(x+12)^2 + 839$ • ³ ✓ 1 |
| <p>Candidate E</p> $-6(x-2)^2 - 1$ Check: $= -6(x^2 - 4x + 4) - 1$ $= -6x^2 + 24x - 24 - 1$ $= -6x^2 + 24x - 25$ Award 3/3 | <p>Candidate F</p> $-6x^2 + 24x - 25$ $= 6x^2 - 24x + 25$ • ¹ ✗ $= 6(x^2 - 4x...)$ $= 6(x-2)^2 ...$ • ² ✓ 1 $= -6(x-2)^2 ...$ • ³ ✗ |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|-----|---|---|----------|
| | (b) | <p>Method 1</p> <ul style="list-style-type: none"> •⁴ differentiate •⁵ link with (a) and identify sign of $(x-2)^2$ •⁶ communicate reason | <p>Method 1</p> <ul style="list-style-type: none"> •⁴ $-6x^2 + 24x - 25$ •⁵ $f'(x) = -6(x-2)^2 - 1$ and $(x-2)^2 \geq 0 \forall x$ •⁶ eg $\therefore -6(x-2)^2 - 1 < 0 \forall x$ \Rightarrow always strictly decreasing | 3 |
| | | <p>Method 2</p> <ul style="list-style-type: none"> •⁴ differentiate •⁵ identify maximum value of $f'(x)$ •⁶ communicate reason | <p>Method 2</p> <ul style="list-style-type: none"> •⁴ $-6x^2 + 24x - 25$ •⁵ 'maximum value is -1' or annotated sketch including x-axis •⁶ $-1 < 0$ or 'graph lies below x-axis' $\therefore f'(x) < 0 \forall x$ \Rightarrow always strictly decreasing | |

Notes:

3. In Method 1, do not penalise $(x-2)^2 > 0$ or the omission of $f'(x)$ at •⁵.
4. In Method 1, accept $-6(x-2)^2 \leq 0$ or $-6(x-2)^2 < 0$ at •⁵.
5. At •⁵ communication must be explicitly in terms of the derivative of the given function. Do not accept statements such as ' $(\text{something})^2 \geq 0$ ', ' $\text{something squared} \geq 0$ '. However, •⁶ is still available.

Commonly Observed Responses:

| | |
|-----------------------------------|------------------|
| Candidate G | |
| $f'(x) = -6x^2 + 24x - 25$ | • ⁴ ✓ |
| $f'(x) = -6(x-2)^2 - 1$ | • ⁵ ^ |
| $-6(x-2)^2 - 1 < 0$ | |
| \Rightarrow strictly decreasing | • ⁶ ^ |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|--|-----|--|--|----------|
| 8. | (a) | <p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ equate composite function to x •² write $f(f^{-1}(x))$ in terms of $f^{-1}(x)$ •³ state inverse function | <p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ $f(f^{-1}(x)) = x$ •² $\sqrt[3]{f^{-1}(x)} + 8 = x$ •³ $f^{-1}(x) = (x-8)^3$ | 3 |
| | | <p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ write as $y = f(x)$ and start to rearrange •² express x in terms of y •³ state inverse function | <p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ $y = f(x) \Rightarrow x = f^{-1}(y)$ $y - 8 = \sqrt[3]{x}$ •² $x = (y-8)^3$ •³ $f^{-1}(y) = (y-8)^3$ $\Rightarrow f^{-1}(x) = (x-8)^3$ | |
| Notes: | | | | |
| <ol style="list-style-type: none"> 1. In Method 2, accept '$y - 8 = \sqrt[3]{x}$' without reference to $y = f(x) \Rightarrow x = f^{-1}(y)$ at •¹. 2. In Method 2, accept $f^{-1}(x) = (x-8)^3$ without reference to $f^{-1}(y)$ at •³. 3. At •³ stage, accept f^{-1} written in terms of any dummy variable eg $f^{-1}(y) = (y-8)^3$. 4. $y = (x-8)^3$ does not gain •³. 5. $f^{-1}(x) = (x-8)^3$ with no working gains 3/3. | | | | |

| Question | Generic scheme | Illustrative scheme | Max mark | | | |
|---|-----------------------------|--|-----------|--|---|--|
| Commonly Observed Responses: | | | | | | |
| Candidate A - multiple expressions for $y = f(x)$ $f(x) = \sqrt[3]{x} + 8$ $y = \sqrt[3]{x} + 8$ $y - 8 = \sqrt[3]{x}$ $x = (y - 8)^3$ $y = (x - 8)^3$ $f^{-1}(x) = (x - 8)^3$ | Award 2/3 | Candidate B - multiple expressions for $y = f(x)$ $f(x) = \sqrt[3]{x} + 8$ $y = \sqrt[3]{x} + 8$ $x = \sqrt[3]{y} + 8$ $y = (x - 8)^3$ $f^{-1}(x) = (x - 8)^3$ | Award 2/3 | | | |
| Candidate C - BEWARE $f'(x) = \dots$ | • ³ ✗ | Candidate D $f^{-1}(x) = x - 8^3$ with no working | Award 0/3 | | | |
| Candidate E $x \rightarrow \sqrt[3]{x} \rightarrow \sqrt[3]{x} + 8 = f(x)$ $\sqrt[3]{} \rightarrow +8$ $\therefore -8 \rightarrow ()^3$ $(x - 8)^3$ $f^{-1}(x) = (x - 8)^3$ | | | | • ¹ ✓ • ² ✓ • ³ ✓ | <div style="border: 1px solid black; border-radius: 15px; padding: 10px; display: inline-block;"> awarded for knowing to perform inverse operations in reverse </div> | |
| (b) | • ⁴ state domain | • ⁴ $9 \leq x \leq 18, x \in \mathbb{R}$ | 1 | | | |
| Notes: | | | | | | |
| 1. Do not penalise the omission of $x \in \mathbb{R}$. | | | | | | |
| Commonly Observed Responses: | | | | | | |
| | | | | | | |

| Question | | Generic scheme | Illustrative scheme | Max mark | |
|--|-----|--|--|----------|--|
| 9. | (a) | • ¹ identify initial power | • ¹ 120 | 1 | |
| Notes: | | | | | |
| Commonly Observed Responses: | | | | | |
| | (b) | • ² interpret information • ³ process equation • ⁴ write in logarithmic form • ⁵ process for t | • ² $102 = 120e^{-0.0079t}$ stated or implied by • ³ • ³ $e^{-0.0079t} = 0.85$ • ⁴ $\log_e 0.85 = -0.0079t$ • ⁵ 20.572... | 4 | |
| Notes: | | | | | |
| 1. Candidates who interpret 15% incorrectly do not gain • ² , but • ³ , • ⁴ and • ⁵ are still available. See Candidate E. 2. • ³ may be implied by • ⁴ . 3. Any base may be used at • ⁴ stage. See Candidate A. 4. Accept $\ln 0.85 = -0.0079t \ln e$ for • ⁴ . 5. Accept 20.57 or 20.6 at • ⁵ . 6. The calculation at • ⁵ must follow from the valid use of exponentials and logarithms at • ³ and • ⁴ . 7. For candidates who take an iterative approach to arrive at $t = 20.6$ award 1/4. However, if, in the iterations P_t is evaluated for $t = 20.55$ and $t = 20.65$ then award 4/4. | | | | | |
| Commonly Observed Responses: | | | | | |
| Candidate A | | $102 = 120e^{-0.0079t}$ • ² ✓ $e^{-0.0079t} = 0.85$ • ³ ✓ $\log_{10} 0.85 = -0.0079t \log_{10} e$ • ⁴ ✓ 20.6 • ⁵ ✓ | Candidate B | | |
| Candidate C | | $\log_e 0.85 = -0.0079t$ • ⁴ ✓ $t = 20.6$ years • ⁵ ✓ $t = 20$ years 6 months | $102 = 120e^{-0.0079t}$ • ² ✓ $e^{-0.0079t} = 0.85$ • ³ ✓ $t = 20.6$ • ⁴ ^ • ⁵ ✓ 1 | | |
| Candidate D | | $\log_e 0.85 = -0.0079t$ • ⁴ ✓ $t = 20$ years 6 months • ⁵ ✗ | | | |
| Candidate E | | $15 = 100e^{-0.0079t}$ $e^{-0.0079t} = 0.15$ • ³ ✓ 1 $\log_e 0.15 = -0.0079t$ • ⁴ ✓ 1 240.1... • ⁵ ✓ 1 | Incorrect conversion subsequent to answer is not penalised | | |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|-----|--|--|----------|
| 10. | (a) | <ul style="list-style-type: none"> •¹ use -3 in synthetic division or in evaluation of quartic •² complete division/evaluation and interpret result | <ul style="list-style-type: none"> •¹ $\begin{array}{r rrrrr} -3 & 3 & 10 & 1 & -8 & -6 \\ & & & & & 3 \end{array}$ or $3 \times (-3)^4 + 10 \times (-3)^3 + (-3)^2 - 8 \times (-3) - 6$ •² $\begin{array}{r rrrrr} -3 & 3 & 10 & 1 & -8 & -6 \\ & & -9 & -3 & 6 & 6 \\ \hline & 3 & 1 & -2 & -2 & 0 \end{array}$ Remainder = 0 $\therefore (x+3)$ is a factor or $f(-3) = 0 \therefore (x+3)$ is a factor | 2 |

Notes:

1. Communication at •² must be consistent with working at that stage ie a candidate's working must arrive legitimately at 0 before •² can be awarded.
2. Accept any of the following for •²:
 - ' $f(-3) = 0$ so $(x+3)$ is a factor'
 - 'since remainder = 0, it is a factor'
 - the '0' from any method linked to the word 'factor' by 'so', 'hence', \therefore , \rightarrow , \Rightarrow etc.
3. Do not accept any of the following for •²:
 - double underlining the '0' or boxing the '0' without comment
 - ' $x = -3$ is a factor', ' \dots is a root'
 - the word 'factor' only, with no link.

Commonly Observed Responses:

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|--|--|--|----------|
| (b) | | <ul style="list-style-type: none"> •³ identify cubic and attempt to factorise •⁴ find second factor •⁵ identify quadratic •⁶ evaluate discriminant •⁷ interpret discriminant and factorise fully | <ul style="list-style-type: none"> •³ eg $\begin{array}{r rrrr} & 3 & 1 & -2 & -2 \\ & & \dots & \dots & \dots \\ & \dots & \dots & \dots & \dots \end{array}$ •⁴ eg $\begin{array}{r rrrr} 1 & 3 & 1 & -2 & -2 \\ & 3 & 4 & 2 & \\ \hline & 3 & 4 & 2 & 0 \end{array}$ leading to $(x-1)$ •⁵ $3x^2 + 4x + 2$ •⁶ -8 •⁷ since $-8 < 0$, quadratic has no (real) factors leading to $(x+3)(x-1)(3x^2 + 4x + 2)$ | 5 |

Notes:

4. Candidates who arrive at $(x+3)(x-1)(3x^2 + 4x + 2)$ by using algebraic long division or by inspection gain •³, •⁴ and •⁵.
5. Evidence for •⁶ may appear in the quadratic formula.
6. Accept ' $-8 < 0$ so no real roots' with the fully factorised quartic for •⁷:
7. Do not accept any of the following for •⁷:
 - $(x+3)(x-1)(3x^2 + 4x + 2)$ does not factorise
 - $(x+3)(x-1)(\dots \dots)(\dots \dots)$ cannot factorise further.
8. Accept $(x+3)(x-1)3x^2 + 4x + 2$, with a valid reason for •⁷.
9. Where the quadratic factor obtained at •⁵ can be factorised, •⁶ and •⁷ are not available.

Commonly Observed Responses:

| Candidate A | | Candidate B | |
|-----------------------------|--|-----------------------------|------------------|
| $(x+3)(x-1)(3x^2 + 4x + 2)$ | • ⁵ ✓ | $(x+3)(x-1)(3x^2 + 4x + 2)$ | • ⁵ ✓ |
| $b^2 - 4ac = 16 - 24 < 0$ | • ⁶ ^ | $b^2 - 4ac < 0$ | • ⁶ ^ |
| so does not factorise | • ⁷ ✓ 1 | so does not factorise | • ⁷ ^ |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|--|-----|---|---|----------|
| 11. | (a) | <ul style="list-style-type: none"> •¹ express A in terms of x and h •² express height in terms of x •³ substitute for h and complete proof | <ul style="list-style-type: none"> •¹ $(A=)16x^2 + 16xh$ •² $h = \frac{2000}{8x^2}$ •³ $A = 16x^2 + 16x \times \frac{2000}{8x^2}$ leading to $A = 16x^2 + \frac{4000}{x}$ | 3 |
| Notes: | | | | |
| <ol style="list-style-type: none"> 1. At •¹ accept any unsimplified form of $16x^2 + 16xh$. 2. The substitution for h at •³ must be clearly shown for •³ to be available. 3. For candidates who omit some of the surfaces of the box, only •² is available. | | | | |
| Commonly Observed Responses: | | | | |
| | (b) | <ul style="list-style-type: none"> •⁴ express A in differentiable form •⁵ differentiate •⁶ equate expression for derivative to 0 •⁷ process for x •⁸ verify nature •⁹ evaluate A | <ul style="list-style-type: none"> •⁴ $16x^2 + 4000x^{-1}$ •⁵ $32x - 4000x^{-2}$ •⁶ $32x - 4000x^{-2} = 0$ •⁷ 5 •⁸ table of signs for a derivative (see below) \therefore minimum or $A''(x) = 96 > 0 \Rightarrow$ minimum •⁹ $A = 1200$ or min value = 1200 | 6 |

Notes:

4. For a numerical approach award 0/6.
5. •⁶ can be awarded for $32x = 4000x^{-2}$.
6. For candidates who integrate any term at the •⁵ stage, only •⁶ is available on follow through for setting their 'derivative' to 0.
7. •⁷, •⁸ and •⁹ are only available for working with a derivative which contains an index ≤ -2 .
8. $\sqrt[3]{\frac{4000}{32}}$ must be simplified at •⁷ or •⁸ for •⁷ to be awarded.
9. •⁸ is not available to candidates who consider a value of $x \leq 0$ in the neighbourhood of 5.
10. •⁹ is still available in cases where a candidate's table of signs does not lead legitimately to a minimum at •⁸.
11. •⁸ and •⁹ are not available to candidates who state that the minimum exists at a negative value of x . See Candidates C and D.

For the table of signs for a derivative, accept:

| | | | |
|----------------|-------|-----|-------|
| x | 5^- | 5 | 5^+ |
| $A'(x)$ | - | 0 | + |
| Shape or slope | \ | - | / |

| | | | |
|----------------|---------------|-----|---------------|
| x | \rightarrow | 5 | \rightarrow |
| $A'(x)$ | - | 0 | + |
| Shape or slope | \ | - | / |

| | | | |
|----------------|-----|-----|-----|
| x | a | 5 | b |
| $A'(x)$ | - | 0 | + |
| Shape or slope | \ | - | / |

Arrows are taken to mean 'in the neighbourhood of'

Where $0 < a < 5$ and $b > 5$

- For this question do not penalise the omission of 'x' or the word 'shape'/'slope'.
- Stating values of $A'(x)$ in the table is an acceptable alternative to writing '+' or '-' signs. Values must be checked for accuracy.
- The only acceptable variations of $A'(x)$ are: A' , $\frac{dA}{dx}$ and $32x - 4000x^{-2}$.

Commonly Observed Responses:

Candidate A - differentiating over multiple lines

$A'(x) = 32x + 4000x^{-1}$ •⁴ ^
 $A'(x) = 32x - 4000x^{-2}$ •⁵ x
 $32x - 4000x^{-2} = 0$ •⁶ ✓ 1

Candidate B - differentiating over multiple lines

$A = 16x^2 + 4000x^{-1}$ •⁴ ✓
 $A'(x) = 32x + 4000x^{-1}$
 $A'(x) = 32x - 4000x^{-2}$ •⁵ x
 $32x - 4000x^{-2} = 0$ •⁶ ✓ 1

Candidate C - only considers 5

$A = 16x^2 + 4000x^{-1}$ •⁴ ✓
 $A' = 32x - 4000x^{-2} = 0$ •⁵ ✓ •⁶ ✓
 $x = \pm 5$ •⁷ x

| | | | |
|------|---------------|-----|---------------|
| x | \rightarrow | 5 | \rightarrow |
| A' | - | 0 | + |
| | \ | - | / |

 \therefore minimum •⁸ ✓ 1
 $A = 1200$ or min value = 1200 •⁹ ✓ 1

Candidate D - considers 5 and negative 5 in separate tables

$A = 16x^2 + 4000x^{-1}$ •⁴ ✓
 $A' = 32x - 4000x^{-2} = 0$ •⁵ ✓ •⁶ ✓
 $x = \pm 5$ •⁷ x

| | | | |
|------|---------------|-----|---------------|
| x | \rightarrow | 5 | \rightarrow |
| A' | - | 0 | + |
| | \ | - | / |

| | | | |
|------|---------------|------|---------------|
| x | \rightarrow | -5 | \rightarrow |
| A' | - | 0 | + |
| | / | - | \ |

 \therefore minimum when $x = 5$ •⁸ ✓ 1
 $A = 1200$ or min value = 1200 •⁹ ✓ 1

Ignore incorrect working in second table

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|--|--|---|----------|
| 12. | | <p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ state linear equation •² introduce logs •³ use laws of logs •⁴ use laws of logs •⁵ state a and b | <p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ $\log_4 y = 3x - 1$ •² $\log_4 y = 3x \log_4 4 - \log_4 4$ •³ $\log_4 y = \log_4 4^{3x} - \log_4 4$ •⁴ $\log_4 y = \log_4 \left(\frac{4^{3x}}{4} \right)$ or $\log_4 y = \log_4 4^{-1} 4^{3x}$ •⁵ $a = \frac{1}{4}, b = 64$ | 5 |
| | | <p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ state linear equation •² convert to exponential form •³ use laws of indices •⁴ state a •⁵ state b | <p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ $\log_4 y = 3x - 1$ •² $y = 4^{3x-1}$ •³ $y = 4^{-1} 4^{3x}$ •⁴ $a = \frac{1}{4}$ •⁵ $b = 64$ | 5 |
| | | <p style="text-align: center;">Method 3</p> <ul style="list-style-type: none"> •¹ introduce logs to $y = ab^x$ •² use laws of logs •³ interpret intercept •⁴ interpret gradient •⁵ state a and b | <p style="text-align: center;">Method 3</p> <p style="text-align: center;">The equations at •¹, •², •³ and •⁴ must be stated explicitly.</p> <ul style="list-style-type: none"> •¹ $\log_4 y = \log_4 ab^x$ •² $\log_4 y = \log_4 a + x \log_4 b$ •³ $-1 = \log_4 a$ •⁴ $3 = \log_4 b$ •⁵ $a = \frac{1}{4}, b = 64$ | 5 |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|---|--|--|--|----------|
| | | <p style="text-align: center;">Method 4</p> <ul style="list-style-type: none"> •¹ interpret point on log graph •² convert from log to exponential form •³ interpret point and convert •⁴ substitute into $y = ab^x$ and evaluate a •⁵ substitute other point into $y = ab^x$ and evaluate b | <p style="text-align: center;">Method 4</p> <ul style="list-style-type: none"> •¹ $x = 3$ and $\log_4 y = 8$ •² $x = 3$ and $y = 4^8$ •³ $x = 0$ and $\log_4 y = -1$ $x = 0$ and $y = 4^{-1}$ •⁴ $4^{-1} = ab^0 \Rightarrow a = \frac{1}{4}$ •⁵ $4^8 = \frac{1}{4}b^3 \Rightarrow b = 64$ | 5 |
| Notes: | | | | |
| <ol style="list-style-type: none"> 1. In any method, marks may only be awarded within a valid strategy using $y = ab^x$. 2. Accept $y = \frac{1}{4} \cdot 64^x$ for •⁵. 3. Markers must identify the method which best matches the candidates approach; they must not mix and match between methods. 4. Penalise the omission of base 4 at most once in any method. 5. Do not accept $a = 4^{-1}$. | | | | |
| Commonly Observed Responses: | | | | |
| | | | | |

| Question | | Generic scheme | Illustrative scheme | Max mark |
|----------|--|---|---|----------|
| 13. | | <ul style="list-style-type: none"> •¹ interpret information given •² integrate any two terms •³ complete integration •⁴ interpret information given and substitute •⁵ process for c and state expression for $f(x)$ | <ul style="list-style-type: none"> •¹ $f'(x) = 3x^2 - 16x + 11$ or $f(x) = \int(3x^2 - 16x + 11)dx$ •² eg $\frac{3x^3}{3} - \frac{16x^2}{2} \dots$ •³ $\dots + 11x + c$ •⁴ $0 = 7^3 - 8 \times 7^2 + 11 \times 7 + c$ •⁵ $f(x) = x^3 - 8x^2 + 11x - 28$ | 5 |

Notes:

1. For candidates who make no attempt to integrate to find $f(x)$ award 0/5.
2. Do not penalise the omission of $f(x)$ or dx or the appearance of $+c$ at •¹.
3. If any two terms have been integrated correctly •¹ may be implied by •².
4. For candidates who omit $+c$, only •¹ and •² are available.
5. For candidates who differentiate **any** term, •³ •⁴ and •⁵ are not available.
6. Candidates must attempt to integrate both terms containing x for •⁴ and •⁵ to be available. See Candidate B.
7. Accept $y = x^3 - 8x^2 + 11x - 28$ at •⁵ since $y = f(x)$ is defined in the question.
8. Candidates must simplify coefficients in **their** final line of working for the last mark available in that line of working to be awarded.

Commonly Observed Responses:

Candidate A - incomplete substitution

$$f(x) = x^3 - 8x^2 + 11x + c \quad \bullet^1 \checkmark \quad \bullet^2 \checkmark \quad \bullet^3 \checkmark$$

$$f(x) = 7^3 - 8 \times 7^2 + 11 \times 7 + c \quad \bullet^4 \wedge$$

$$c = -28$$

$$f(x) = x^3 - 8x^2 + 11x - 28 \quad \bullet^5 \boxed{\checkmark 1}$$

Candidate B - partial integration

$$f(x) = x^3 - 8x^2 + 11x + c \quad \bullet^1 \checkmark \quad \bullet^2 \checkmark \quad \bullet^3 \times$$

$$0 = 7^3 - 8 \times 7^2 + 11 + c \quad \bullet^4 \boxed{\checkmark 1}$$

$$c = 38$$

$$f(x) = x^3 - 8x^2 + 49 \quad \bullet^5 \boxed{\checkmark 1}$$

| Question | Generic scheme | Illustrative scheme | Max mark |
|----------|---|--|----------|
| 14. | <ul style="list-style-type: none"> •¹ expand •² evaluate $\mathbf{u} \cdot \mathbf{u}$ •³ determine equation in $\cos \theta$ •⁴ evaluate angle | <ul style="list-style-type: none"> •¹ $\mathbf{u} \cdot \mathbf{u} + \mathbf{u} \cdot \mathbf{v}$ •² 16 •³ $20 \cos \theta = 5$ or $\cos \theta = \frac{5}{20}$ •⁴ $75 \cdot 5 \dots^\circ$ or $1 \cdot 31 \dots$ radians | 4 |

Notes:

1. Do not accept \mathbf{u}^2 for •¹, however •², •³ and •⁴ are still available.
2. Where there is no evidence for •¹, then •², •³ and •⁴ are not available, however see Candidates C and D.
3. Where candidates use $|\mathbf{u}| \neq 4$, then •³ and •⁴ are not available.
4. Where there is no evidence of using $|\mathbf{u}|^2$, •³ is not available. See Candidate A.
5. Do not penalise omission of units in final answer.
6. Ignore the appearance of $284 \cdot 5^\circ$.
7. Accept answers which round to 76° or 1.3 radians.

Commonly Observed Responses:

| | | | |
|--|--|--|---|
| <p>Candidate A</p> $\mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) = \mathbf{u} \cdot \mathbf{u} + \mathbf{u} \cdot \mathbf{v}$ $4 + 20 \cos \theta = 21$ $\cos \theta = \frac{17}{20}$ $\theta = 31 \cdot 7 \dots^\circ$ | <ul style="list-style-type: none"> •¹ ✓ •² ✗ •³ ✓ 2 •⁴ ✓ 1 | <p>Candidate B</p> $16 + \mathbf{u} \cdot \mathbf{v} = 21$ $\mathbf{u} \cdot \mathbf{v} = 5$ $\cos \theta = \frac{5}{20}$ $\theta = 75 \cdot 5^\circ$ | <ul style="list-style-type: none"> •¹ ✓ •² ✓ •³ ✓ •⁴ ✓ |
| <p>Candidate C - missing working</p> $\mathbf{u} \cdot \mathbf{u} = 16$ $\mathbf{u} \cdot \mathbf{v} = 21 - 16$ $\cos \theta = \frac{5}{20}$ $\theta = 75 \cdot 5^\circ$ | <ul style="list-style-type: none"> •² ✓ •¹ ✓ •³ ✓ •⁴ ✓ | <p>Candidate D - missing working</p> $21 - 16 = 5$ $\cos \theta = \frac{5}{20}$ $\theta = 75 \cdot 5^\circ$ | <ul style="list-style-type: none"> •¹ ^ •² ✓ •³ ✓ •⁴ ✓ |

| Question | | Generic scheme | Illustrative scheme | Max mark | |
|---|-----|---|--|---|---|
| 15. | (a) | <ul style="list-style-type: none"> •¹ find gradient of radius •² state gradient of tangent •³ state equation of tangent | <ul style="list-style-type: none"> •¹ $-\frac{1}{3}$ •² 3 •³ $y = 3x - 2$ | 3 | |
| Notes: | | | | | |
| <p>1. Do not accept $y = \frac{3}{1}x - 2$ for •³.</p> <p>2. •³ is only available as a consequence of trying to find and use a perpendicular gradient.</p> <p>3. At •³ accept, $y - 3x + 2 = 0$ or any other rearrangement of the equation where the constant terms have been simplified.</p> | | | | | |
| Commonly Observed Responses: | | | | | |
| | (b) | (i) | • ⁴ find coordinates of T | • ⁴ (0, -2) | 1 |
| | | (ii) | <ul style="list-style-type: none"> •⁵ find midpoint CT •⁶ find radius of circle with diameter CT •⁷ state equation of circle | <ul style="list-style-type: none"> •⁵ (4,5) •⁶ $\sqrt{65}$ stated or implied by •⁷ •⁷ $(x-4)^2 + (y-5)^2 = 65$ | 3 |
| Notes: | | | | | |
| <p>4. Answers in part (b)(i) must be consistent with answers from part (a).</p> <p>5. Accept $x = 0, y = -2$ for •⁴.</p> <p>6. $(x-4)^2 + (y-5)^2 = (\sqrt{65})^2$ does not gain •⁷.</p> <p>7. •⁷ is not available to candidates who use a line other than CT as the diameter of the circle.</p> | | | | | |
| Commonly Observed Responses: | | | | | |

[END OF MARKING INSTRUCTIONS]